

In collaboration with  
Róbert Vértési

# CORRELATION OF HEAVY AND LIGHT FLAVOURS IN SIMULATIONS

Eszter Frajna

BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS

18th Zimányi School - Winter Workshop on Heavy Ion Physics

2018. December 4.

# Fundamentals of high-energy physics

## Standard Model of Elementary Particles


		three generations of matter (fermions)			interactions / force carriers (bosons)	
		I	II	III		
mass		$\approx 2.2 \text{ MeV}/c^2$	$\approx 1.28 \text{ GeV}/c^2$	$\approx 173.1 \text{ GeV}/c^2$	0	$\approx 125.09 \text{ GeV}/c^2$
charge	$2/3$	$2/3$	$2/3$	$2/3$	0	0
spin	$1/2$	$1/2$	$1/2$	$1/2$	1	0
		<b>U</b> up	<b>C</b> charm	<b>T</b> top	<b>g</b> gluon	<b>H</b> higgs
		<b>D</b> down	<b>S</b> strange	<b>B</b> bottom	<b><math>\gamma</math></b> photon	
		<b>E</b> electron	<b><math>\mu</math></b> muon	<b><math>\tau</math></b> tau	<b>Z</b> Z boson	
		<b><math>\nu_e</math></b> electron neutrino	<b><math>\nu_\mu</math></b> muon neutrino	<b><math>\nu_\tau</math></b> tau neutrino	<b>W</b> W boson	

**QUARKS** (left side of fermion table)


**LEPTONS** (left side of fermion table)

**GAUGE BOSONS VECTOR BOSONS** (left side of boson table)

**SCALAR BOSONS** (right side of boson table)




- QGP can be recreated in the lab in heavy ion collisions



It was the first convincing evidence of hot and dense strongly interacting nuclear matter in the final state.

Based on the angular correlation data of STAR, jet-quenching can be observed.



## Previous measurements in heavy ion collisions

Variables determining the particle movement:

- ❖ Rapidity

$$y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z}$$

- ❖ Pseudorapidity

$$\eta = \frac{1}{2} \ln \frac{p + p_z}{p - p_z} = -\ln \left( \tanh \left( \frac{\theta}{2} \right) \right)$$

- ❖ Transverse momentum



$$p_T = \sqrt{p_x^2 + p_y^2}$$

- ❖ Azimuthal angle

$$\Phi = \frac{p_y}{p_x}$$

# Two-particle correlations


- select a trigger particle from a given momentum (pT) range
- then in a lower momentum (pT) window examine all the other particles from the same event (associated particles)
- near-side and away-side peak



# Simulation settings

- Event generator based on Monte Carlo method: PYTHIA 8.1
- Hard QCD events were created in PYTHIA using the default Monash 2013 settings for LHC p + p data.
- The phase space has been reduced so that the leading hard process has at least 5 GeV / c momentum.
- In the case of heavy quark samples, the renormalization scales were set based on the STAR Heavy Flavor tune (from which we expect the corresponding result in the examined momentum range) and only the following two processes are allowed:  $gg \rightarrow b\bar{b}$ ,  $qq \rightarrow b\bar{b}$  and  $gg \rightarrow c\bar{c}$ ,  $qq \rightarrow c\bar{c}$ .

# Acceptance correction



$$\frac{dN}{d\Delta\eta} = \frac{1}{2A} - \frac{1}{4A^2} |\Delta\eta|$$

To correct the finite size of the detector we divide the correlation in the  $\Delta\eta - \Delta\phi$  plane with this tent-shaped function.

# Applications of correlation measurements

## ❖ Understanding jet structure by correlation

- Interaction of partons with QGP can be studied by full jet reconstruction
- The background size makes it difficult to reconstruct the jet under a certain momentum
- Solution: measuring the angular correlation of particles

## ❖ Identify the characteristic correlation images of heavy quarks

- Compare near- and away-side peaks associated with hadrons from different heavy quarks from a given  $p_T$
- Which probes are sensitive to heavy quarks?

## ❖ Parameterization of correlation images by fitting

# Determining the peak's parameters by function fitting

Gauss

$$N \cdot \frac{1}{\sqrt{2\pi} \sigma_\eta} \cdot e^{-\left(\frac{\Delta\eta^2}{2\sigma_\eta^2}\right)}$$

$$N \cdot \frac{1}{\sqrt{2\pi} \sigma_\Phi} \cdot e^{-\left(\frac{\Delta\eta^2}{2\sigma_\Phi^2}\right)}$$

Generalized Gauss

*If  $\gamma = 2$ , it can be identified with a simple gaussian function*


$$N \cdot \frac{\gamma_\eta}{2\omega_\eta \Gamma\left(\frac{1}{\gamma_\eta}\right)} \cdot e^{-\left(\frac{|\Delta\eta|}{\omega_\eta}\right)^{\gamma_\eta}}$$

$$N \cdot \frac{\gamma_\Phi}{2\omega_\Phi \Gamma\left(\frac{1}{\gamma_\Phi}\right)} \cdot e^{-\left(\frac{|\Delta\Phi|}{\omega_\Phi}\right)^{\gamma_\Phi}}$$




# RESULTS

# Correlations of light charged hadrons





The near-side peak is significantly "peakier" than a Gaussian,  $\gamma \sim 1.016$ .



The near-side and away-side peaks in  $\Delta\phi$  are well described by a Gaussian


# Correlations of light charged hadrons




Peaks are getting narrower towards higher  $p_T$  (Lorentz-boost).  
GenGauss parameter  $\gamma$  is constant within error.

# Prompt production of heavy flavour mesons

D-meson from c-quarks




B-meson from b-quarks




Peaks are consistent with Gaussian

# D-meson from the decay of the B-meson





Strong dependence of away-side peak on  $pT$  (not observed in light flavor or prompt heavy flavour production)



GenGauss parameter  $\gamma$  decreases with  $pT$ , together with  $\sigma$ . (Peaks are getting both narrower and "peakier" towards high  $pT$ )

# Investigation of electrons from B-mesons





Wide correlation pictures compared to hadrons.  
No significant dependence of  $\gamma$  on  $p_T$ .

# Comparison of B-meson and b-quark

b-quark

B-meson



Evolution of correlation pictures match within uncertainties.  
B meson is a good proxy for the b quark.

# SUMMARY

- ❖ The shape of the correlation peaks can be used to separate the electrons coming from b-quark decays. This could be a method of identification that, combined with ITS identification, may provide a much better sample purity for examining the secondary vertex shift.
- ❖ Correlation images are sensitive to the distribution of secondary vertex in different quarks, and these processes can be statistically separated from light quarks.
- ❖ It is possible to distinguish which D-meson comes directly and which later decay. This allows the measurement to be used for statistical separation of prompt and late D-mesons.
- ❖ We also have a characteristic b-correlation image, which is present in both b-quarks and B-mesons. B-mesons can be used to study b-quarks.



**THANKS FOR YOUR  
ATTENTION!**