

Tsallis-Thermometer as a QGP Indicator for Large & Small Collisional Systems

Gergely Gábor Barnaföldi¹, Gábor Bíró¹², Tamás Sándor Bíró¹, Aditya Nath Mishra³, Guy Paic⁴

¹Wigner Research Centre for Physics, 29-33 Konkoly-Thege Miklós street, 1121, Budapest, Hungary

²Institute of Physics, Eötvös Loránd University, 1/A Pázmány Péter Sétány, H-1117 Budapest, Hungary.

³University Centre For Research and Development (UCRD), Chandigarh University, Mohali-140413, Punjab, India

⁴Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Apartado Postal 70-543, México City 04510, México




Introduction

Recent experimental results present collectivity also in small systems with high-multiplicity. Today these phenomena are not completely understood: it is an important question whether the presence of the QGP is necessary for the observed collectivity or not. Moreover, the connection between the experimental observables and theories is not trivial. In our phenomenological study we introduce the 'Tsallis-thermometer' as an indicator of QGP, that result in the description of the smooth transition from small to large collisional systems. The method also works well with geometry-selection of an event and correlates well with sphericity classified events. Results enable us to qualitatively define the underlying event definition beyond the CDF definition.

Tsallis distributions

Thermodynamically consistent fit function:

$$\frac{d^2N}{2\pi p_T dp_T dy} \Big|_{y \approx 0} = Am_T \left[1 + \frac{q-1}{T}(m_T - m) \right]^{-\frac{q}{q-1}}$$



Investigated experimental data:


$\pi^\pm, K^\pm, K_s^0, K^{*0}, p(\bar{p}), \Phi, \Lambda^0, \Xi^\pm, \Sigma^\pm, \Xi^0, \Omega^\pm$

$\sqrt{s_{NN}} \in 130 \text{ GeV}, 13 \text{ TeV}$

pp, p-A and AA collisions

Various multiplicity classes


Consistency: $P = T_s + \mu n - \varepsilon$



Scaling with UE activity



Underlying event activity can be quantitatively seen by the Tsallis thermometer. Geometrical scanning is correlates with sphericity classified events.



Acknowledgement

This work was supported by the Hungarian National Research Fund OTKA grant K135515, NKFIH 2019-2.1.1-TÉT-2019-00078, 2019-2.1.1-TÉT-2019-00050; Wigner Scientific Computing Laboratory (WSCLAB) (the former Wigner GPU Laboratory), and by the Ministry of Innovation and Technology NRD Office within the framework of the MILAB Artificial Intelligence National Laboratory Program.


Hadron spectra and multiplicity scaling

Scaling with multiplicity & center of mass energy


$$T(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = T_0 + T_1 \ln \frac{\sqrt{s_{NN}}}{m} + T_2 \ln \ln \langle N_{ch}/\eta \rangle,$$

$$q(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = q_0 + q_1 \ln \frac{\sqrt{s_{NN}}}{m} + q_2 \ln \ln \langle N_{ch}/\eta \rangle,$$

Strong correlation between the parameters has been recognized as function of multiplicity:



... GIVES YOU MORE:
→ TEMPERATURE QUANTITY: T
→ INFO ON SYSTEM (SIZE): q





Further correlation with multiplicity fluctuation – assuming NBD:

$$T = E(\delta^2 - (q - 1))$$

Equation of State

EoS can be obtained, due to the thermodynamical consistency:

- Hadron mass order
- Present evolution
- Branches compatible with hadron d.o.f.



Summary

The Tsallis-thermometer is an excellent measure for locating collective effects in high-energy collisions from small to large system sizes. Scaling behaviors are inherited both from the QCD nature and from system size effects. Thermodynamical consistency helps us to measure EoS more accurately.

References

- [1] G Bíró, GG Barnaföldi, TS Bíró, J. Phys. G, 47(10) (2020), 105002.
- [2] G Bíró, GG Barnaföldi, K Ürmössy, TS Bíró, Á. Takács, Entropy, 19(3), (2017), 88
- [3] G Bíró, GG Barnaföldi, TS Bíró, K Shen, EPJ Web Conf., 171, (2018), 14008
- [4] G Bíró, GG Barnaföldi, G Papp, TS Bíró, Universe 5 (2019) no.6, 134
- [5] AN Mishra, GG Barnaföldi, G Paic, J. Phys.G 50 (2023) 9, 095004